

ORIGINAL RESEARCH

The acute effects of foam rolling and dynamic stretching protocols on stroke speed performance in adolescent female tennis players

¹Neslihan Sak  ²Ali Tatlıcı 

¹Selcuk University, Faculty of Sport Sciences, Konya, Türkiye

²Selcuk University, Faculty of Sport Sciences, Konya, Türkiye.

Received:

July 09, 2025

Accepted:

September 20, 2025

Online**Published:**

September 25, 2025

Keywords:

Tennis,
Adolescent
athletes, Foam
rolling,
Dynamic
stretching.

Abstract. The aim of this study was to investigate and compare the acute effects of foam rolling and dynamic stretching protocols on the stroke speed performance (forehand, backhand, and serve) of adolescent female tennis players. Ten adolescent female tennis players aged 10–13 years, with 3–4 years of playing experience, voluntarily participated in the study at a tennis club in Ankara. Participants had a mean age of 11.50 ± 1.08 years, mean height of 150.50 ± 6.12 cm, mean weight of 41.10 ± 6.66 kg, and an average tennis training background of 3.20 ± 0.42 years. After completing a general tennis-specific warm-up, participants were randomly divided into two groups. The first group performed 8 minutes of dynamic stretching, while the second group performed 8 minutes of foam rolling. Stroke speeds (forehand, backhand, and serve) were measured using a radar gun following each protocol. After a 48-hour recovery period, a crossover design was implemented and the protocols were repeated with the groups switched. Data were analyzed using SPSS software, and paired sample t-tests were used for comparisons. Following the dynamic stretching protocol, mean forehand speed was 80.00 ± 8.60 km/h, backhand speed was 75.60 ± 11.84 km/h, and serve speed was 80.10 ± 16.87 km/h. After foam rolling, forehand speed was 78.30 ± 14.05 km/h, backhand speed was 75.00 ± 12.18 km/h, and serve speed was 80.60 ± 15.43 km/h. No statistically significant differences were found between the two protocols in any of the stroke speed measurements ($p > 0.05$). The results indicate that neither foam rolling nor dynamic stretching provides a significant acute advantage in improving stroke speed performance in adolescent female tennis players. However, both methods can be considered effective components of a warm-up routine.

Introduction

Tennis is a dynamic sport that combines speed, agility, endurance, technical, and psychological skills (Kovacs, 2006). The forehand, backhand, and serve are fundamental strokes in tennis, each involving technical and biomechanical components that significantly affect a player's overall performance (Irawan et al., 2023; Kovacs, 2006). Moreover, modern tennis requires players to anticipate opponents' actions, adjust strategies quickly, and maintain consistent strokes, making these skills essential for success (Kovacs, 2006; Taşkuyu & Özcan, 2025). Analyzing each stroke biomechanically is crucial for technical development (Elliott, 2006; Irawan et al., 2023).

Biomechanical analysis serves as an essential tool in understanding tennis stroke velocity. Among the key factors influencing stroke speed

are the athlete's arm motion and body positioning. Proper arm acceleration and an optimal body position during the stroke especially in high-speed actions like the forehand can significantly enhance performance (Irawan et al., 2023). Stroke skill, which is a core component of performance, is directly linked not only to the player's technical proficiency but also to an optimal preparatory routine. Therefore, various training methods and techniques aiming to improve tennis performance are emphasized in tennis performance development (Kovacs, 2007).

To increase stroke speed, tennis players must be able to efficiently utilize all parts of their bodies. In particular, the interaction between the upper body (arms and shoulders) and the lower body (legs and core muscles) is one of the most important biomechanical factors contributing to

✉ N. Sak, e-mail; nesli.sak06@gmail.com

Cite: Sak, N. ve Tatlıcı, A. 2025. The acute effects of foam rolling and dynamic stretching protocols on stroke speed performance in adolescent female tennis players. *Journal of National Kinesiology*. 6(2), 140-145.

the stroke speed (Elliott et al., 2009). Furthermore, strength and flexibility training are essential in maximizing this biomechanical synergy (McGill, 2010). Consequently, pre-match or pre-training warm-up protocols are vital for activating the musculoskeletal system, improving stroke performance, and reducing the risk of injury (Ayala et al., 2016; Behm & Chaouachi, 2011).

Numerous studies have examined the effects of warm-up methods on various athletic performance parameters, comparing static and dynamic stretching protocols (McHugh & Cosgrave, 2010), as well as foam rolling and its combination with warm-up routines (Cheatham et al., 2015).

Dynamic stretching exercises are active movements that involve lengthening the muscles while in motion. These exercises can prepare the musculoskeletal system and support neuromuscular activation of the body (Yamaguchi & Ishii, 2005). Dynamic stretching enables the body to achieve the flexibility and range of motion necessary for optimal performance (Opplert & Babault, 2018). Studies in the literature have shown that dynamic stretching improves athletes' strength and power output during training (Behm & Chaouachi, 2011). It can be considered a widely used and traditional type of stretching.

Recently, foam rolling exercises, often used in dynamic sports, have gained popularity. Foam rolling helps increase range of motion, increases blood flow, and reduces muscle soreness (Cheatham et al., 2015). These benefits can make it a potentially effective method for improving stroke speed in tennis players. Foam rolling facilitates the efficient use of body synergy by helping relax the leg and gluteal muscles (MacDonald et al., 2014). In addition to its performance-enhancing effects, foam rolling has also been shown to accelerate recovery and reduce muscle fatigue after exercise (Schroeder & Best, 2015).

However, despite the growing use of dynamic stretching and foam rolling, there is limited evidence regarding their immediate effects on tennis-specific performance, particularly stroke speed. Moreover, most previous studies have focused on adult athletes, leaving a gap in the literature concerning adolescent female players. Addressing this gap is important because physical development, neuromuscular control, and

recovery capacity may differ in this population. Based on this background, the present study aimed to investigate the acute effects of two different warm-up protocols dynamic stretching and foam rolling on stroke speed performance in adolescent female tennis players.

Material and Method

Participants

The study was conducted with the voluntary participation of 10 adolescent female tennis players (aged 10–13 years) who had 3 to 4 years of tennis experience and were actively training at Tennis Clinic, a tennis club located in Ankara, Turkey. Participants were selected based on the following inclusion criteria: regular engagement in physical activity and tennis training, no known health problems, and no physical limitations. All performance assessments were conducted prior to training sessions to eliminate potential fatigue effects that might influence the results. The demographic characteristics of the participants were as follows: mean age 11.50 ± 1.08 years, mean height 150.50 ± 6.12 cm, mean body weight 41.10 ± 6.66 kg, and mean tennis experience 3.20 ± 0.42 years. All participants were informed about the study's objectives and potential risks. The study was approved by the local ethics committee (Protocol number 167, 12/02/2025, Ethics Committee of Selçuk University, Faculty of Sports Science, Konya, Turkey). Before the assessment, every participant received the same detailed information about the testing procedure. Informed consent forms were signed by all participants and their legal guardians in accordance with ethical research standards.

Study Design

This study examined the acute effects of two different warm-up protocols, dynamic stretching and foam rolling, on stroke speed performance in adolescent female tennis players. A randomized crossover design was employed. In the first session, participants were randomly assigned to two groups. The dynamic stretching group performed exercises targeting the quadriceps, hamstrings, gluteus maximus, and gastrocnemius muscles. Each muscle group was stretched for 60 seconds, with the total duration lasting 8 minutes. Movements were performed in a slow and controlled manner to avoid ballistic actions. The timing of each exercise was based on the procedures reported by (Behara & Jacobson, 2017). The foam rolling group applied foam

rolling to the same muscle groups (quadriceps, hamstrings, gluteus maximus, gastrocnemius). Rolling was performed unilaterally, with each muscle group treated for 60 seconds, totaling 8 minutes of application. As with the stretching protocol, the duration of rolling was aligned with the standardized procedures described by Behara and Jacobson (2017). Immediately after the interventions, stroke speed tests were conducted. Forehand, backhand, and serve speeds were measured using a Busnell radar gun (Sports Radar, Bushnell, USA). The radar device was placed 4 meters behind the athlete by an assistant and angled toward the direction of the ball's exit from the racket (Kara et al., 2015). Each stroke was performed 5 times, and the best attempt was recorded for analysis. To minimize residual effects, a 48-hour washout period was implemented. After this period, the groups switched protocols: participants who had performed dynamic stretching in the first session completed the foam rolling protocol, while those who had performed foam rolling completed the dynamic stretching protocol. The same testing procedures were repeated in the second session. This crossover design allowed each athlete to serve as her own control, thereby improving

reliability of comparisons and reducing inter-individual variability.

Statistical Analysis

All statistical analyses were performed using IBM SPSS Statistics version 25.0 (IBM Corp., Armonk, NY, USA). Descriptive statistics (mean $\pm$ standard deviation) were calculated for all performance variables, including stroke speeds (forehand, backhand, and serve) under both warm-up conditions (dynamic stretching and foam rolling). To evaluate the normality of the data, both Shapiro-Wilk tests and visual inspection of skewness and kurtosis values were conducted. Skewness and kurtosis values falling within the acceptable range of ± 2 were considered indicative of normal distribution. The results confirmed that the data met the assumptions of normality. As each participant completed both intervention protocols in a crossover design, paired samples t-tests were used to compare the stroke speed performances between the two warm-up conditions. The level of statistical significance was set at $p < 0.05$.

Findings

Table 1. Descriptive Characteristics of the Participants

Variable	n	Minimum	Maximum	Mean	SD
Age (years)	10	10	13	11.5	1.08
Height (cm)	10	140	157	150.5	6.12
Weight (kg)	10	30	48	41.1	6.66
Tennis Experience (years)	10	3	4	3.2	0.42

Note. SD = Standard Deviation.

Table 2. Comparison of Participants' Dynamic and Foam Roller Stretching Data

	Mean	Std. Dev.	SE	Confidence interval Lower upper		T	p
Dynamic Forehand	80.00	8.602	1.892	-2.580	5.980	.899	.392
Foam roller Forehand	78.30	14.048					
Dynamic Backhand	75.60	11.843	2.023	-3.977	5.177	.297	.774
Foam roller Backhand	75.00	12.184					
Dynamic Service	80.10	16.868	.910	-2.558	1.558	-.550	.596
Foam roller Service	80.60	15.429					

The average dynamic forehand speed was measured as 80.00 ± 8.602 km/h, while the foam roller forehand speed was 78.30 ± 14.048 km/h. The dynamic backhand speed was found to be

75.60 ± 11.843 km/h, and the foam roller backhand speed was 75.00 ± 12.184 km/h. The dynamic serve speed was recorded as 80.10 ± 16.868 km/h, whereas the foam roller serve

speed was 80.60 ± 15.429 km/h. The t-test analyses revealed that there was no statistically significant difference in stroke speed performance between the dynamic and foam roller warm-up protocols ($p > 0.05$).

Discussion and Conclusion

This study aimed to compare the acute effects of two different warm-up protocols on stroke speed performance in adolescent female tennis players, with the goal of identifying which protocol might be more effective. The findings showed no statistically significant difference between the effects of dynamic and foam roller warm-up protocols on stroke speed performance ($p > 0.05$).

When reviewing similar studies in the literature, it becomes evident that the effects of dynamic stretching and foam roller protocols on muscle performance vary across different sports disciplines (Behm & Chaouachi, 2011; Cheatham et al., 2015). One study reported that dynamic stretching exercises enhance performance, particularly in movements where speed and power are crucial (Fletcher & Jones, 2004). In a tennis-specific study, lower extremity muscle strength was identified as central to producing explosive actions such as the tennis serve (Girard & Millet, 2009). It was also suggested that pre-match stretching might delay the recovery of full muscular performance until near the end of the second set (Knudson et al., 2004). Furthermore, a study reported that a 15-rep dynamic stretching routine before a muscle performance test led to improved performance (Yamaguchi et al., 2007).

Foam roller applications have also shown promise, where increased range of motion may positively affect physical performance (MacDonald et al., 2014). MacDonald and colleagues observed that foam rolling reduced muscle soreness and facilitated recovery following intense exercise. Similarly, a study on basketball players highlighted that vibration foam rolling accelerated short-term recovery and improved physical performance (Li et al., 2023).

In the tennis serve, the “rotator cuff” muscles—supraspinatus for abduction, subscapularis for internal rotation, and infraspinatus and teres minor for external rotation—play a significant role, supported by several other muscles such as pectoralis major and latissimus dorsi (Ozgurbuz & Aksit, 2004; Özgürbüz et al., 2003). These muscle groups

share functional similarities with those used in swimming, where shoulder function and muscular balance are crucial for performance. In fact, Bishop and colleagues have emphasized the importance of shoulder strength and conditioning for sprint swimming performance, particularly in relation to reducing muscular imbalances and improving overall efficiency in the water (Bishop et al., 2013).

In another study comparing dynamic stretching and foam roller warm-up protocols, it was observed that foam roller exercises produced similar results to dynamic stretching in terms of jump and strength performance parameters, while having a favorable impact on joint range of motion (Behara & Jacobson, 2017). Another study on muscle strength reported that foam rolling significantly increased left hamstring strength, whereas dynamic stretching did not produce a significant change (Çakmak, 2021). While many studies indicate that foam rolling enhances muscle elasticity, they also suggest that its effect on performance is not always substantial. The current study is consistent of these findings by demonstrating that both protocols had similar effects on stroke speed.

The measurement results in our study suggest that among tennis players aged 10–13, muscle elasticity and neuromuscular coordination may not be developed enough to create a meaningful difference between the protocols. This could be attributed to the fact that athletes in this age group are still in a developmental phase in terms of muscle structure and flexibility. Another noteworthy finding of the study is that neither protocol had a negative impact on performance. This suggests that coaches can confidently use both dynamic stretching and foam roller routines in various training and pre-match settings.

The findings of this study demonstrated that foam rolling and dynamic stretching protocols did not produce statistically significant acute differences in forehand, backhand, and serve stroke speeds among adolescent female tennis players. However, neither method was significantly inferior to the other. This suggests that both warm-up strategies can be incorporated into pre-competition or pre-training routines by coaches and athletes. While dynamic stretching activates the musculoskeletal system and enhances neuromuscular readiness, foam rolling may improve muscle elasticity and reduce tightness. Thus, both approaches appear to

prepare athletes effectively for tennis-specific performance, even though no clear superiority was observed in acute stroke speed outcomes. Despite these findings, certain limitations of the present study highlight directions for future research. The relatively small sample size restricts the generalizability of the results, and further studies with larger cohorts are recommended to strengthen the evidence. Investigations across different age groups and including male players would also provide a broader understanding of the effects of these interventions. Moreover, as this research focused solely on acute outcomes, examining the long-term effects of dynamic stretching and foam rolling when incorporated into regular training programs would offer valuable insights. Beyond stroke speed, evaluating additional performance parameters such as agility, reaction time, muscular strength, and flexibility could help clarify the broader impact of warm-up protocols. Finally, it would be beneficial to explore whether combining dynamic stretching and foam rolling into a hybrid protocol produces greater benefits compared to applying each method independently.

Financial Support

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of Interest

The authors declare that there is no conflict of interest regarding this article.

Ethics Committee Approval Report

Ethical approval for this study was obtained from the Selçuk University Non-Interventional Clinical Research Ethics Committee Decision (Date: 12.02.2025, Decision No: 167).

Authors' Contribution

Study Design: NS, AT

Data Collection: NS, AT

Statistical Analysis: NS, AT

Manuscript Preparation: NS, AT

Funding Acquisition: NS, AT

References

Ayala, F., Moreno-Pérez, V., Vera-Garcia, F. J., Moya, M., Sanz-Rivas, D., & Fernandez-Fernandez, J. (2016). Acute and time-course effects of

traditional and dynamic warm-up routines in young elite junior tennis players. *PloS one*, 11(4), e0152790.

- Behara, B., & Jacobson, B. H. (2017). Acute effects of deep tissue foam rolling and dynamic stretching on muscular strength, power, and flexibility in division I linemen. *The Journal of Strength & Conditioning Research*, 31(4), 888-892.
- Behm, D. G., & Chaouachi, A. (2011). A review of the acute effects of static and dynamic stretching on performance. *European journal of applied physiology*, 111(11), 2633-2651.
- Bishop, C., Cree, J., Read, P., Chavda, S., Edwards, M., & Turner, A. (2013). Strength and conditioning for sprint swimming. *Strength & Conditioning Journal*, 35(6), 1-6.
- Cheatham, S. W., Kolber, M. J., Cain, M., & Lee, M. (2015). The effects of self-myofascial release using a foam roll or roller massager on joint range of motion, muscle recovery, and performance: a systematic review. *International Journal of Sports Physical Therapy*, 10(6), 827-838.
- Çakmak, D. (2021). *Amatör Voleybol Oyuncularında Statik Germe, Dinamik Germe Ve Foam Roller İle Germenin Performansa Etkisi* [Yüksek lisans, Bahçeşehir Üniversitesi]. İstanbul.
- Elliott, B. (2006). Biomechanics and tennis. *British journal of sports medicine*, 40(5), 392-396.
- Elliott, B., Reid, M., & Crespo, M. (2009). *Technique development in tennis stroke production*. International Tennis Federation.
- Fletcher, I. M., & Jones, B. (2004). The effect of different warm-up stretch protocols on 20 meter sprint performance in trained rugby union players. *Journal of Strength and Conditioning Research*, 18(4), 885-888.
- Girard, O., & Millet, G. P. (2009). Physical determinants of tennis performance in competitive teenage players. *The Journal of Strength & Conditioning Research*, 23(6), 1867-1872.
- Irawan, R., Azam, M., Rahayu, S., Setyawati, H., Soedjatmiko, Adi, S., Priyono, B., & Nugroho, A. (2023). Biomechanical motion of the tennis forehand stroke: Analyzing the impact on the ball speed using biofor analysis software. *Physical Education Theory and Methodology*, 23(6), 918-924.
- Kara, E., Aksit, T., Ozkol, M. Z., & Isık, T. (2015). Effects of 6 week tennis specific exercises program on service velocity. *Turkish Journal of Sport and Exercise*, 17(1), 71-76.
- Knudson, D. V., Noffal, G. J., Bahamonde, R. E., Bauer, J. A., & Blackwell, J. R. (2004). Stretching has no effect on tennis serve performance. *Journal of Strength and Conditioning Research*, 18(3), 654-656.

-
- Kovacs, M. S. (2006). Applied physiology of tennis performance. *British journal of sports medicine*, 40(5), 381-386.
- Kovacs, M. S. (2007). Tennis physiology: training the competitive athlete. *Sports medicine*, 37(3), 189-198.
- Li, F., Song, Y., Cen, X., Sun, D., Lu, Z., Bíró, I., & Gu, Y. (2023). Comparative efficacy of vibration foam rolling and cold water immersion in amateur basketball players after a simulated load of basketball game. *Healthcare*, 11(15), 2178.
- MacDonald, G. Z., Button, D. C., Drinkwater, E. J., & Behm, D. G. (2014). Foam rolling as a recovery tool after an intense bout of physical activity. *Medicine & Science in Sports & Exercise*, 46(1), 131-142.
- McGill, S. (2010). Core training: Evidence translating to better performance and injury prevention. *Strength & Conditioning Journal*, 32(3), 33-46.
- McHugh, M. P., & Cosgrave, C. H. (2010). To stretch or not to stretch: the role of stretching in injury prevention and performance. *Scandinavian journal of medicine & science in sports*, 20(2), 169-181.
- Opplert, J., & Babault, N. (2018). Acute effects of dynamic stretching on muscle flexibility and performance: an analysis of the current literature. *Sports medicine*, 48(2), 299-325.
- Ozgurbuz, C., & Aksit, T. (2004). The concept of a triple-component strength ratio of the shoulder rotator muscles in tennis players. *DEUTSCHE ZEITSCHRIFT FÜR SPORTMEDIZIN*, 55(4), 96-100.
- Özgürbüz, C., Akşit, T., Acar, M. F., Ertat, A., & Durmaz, B. (2003). Elit Tenisçilerde Omuzun Fonksiyonel Kuvvet Ölçümleri. *Spor Hekimliği Dergisi*, 38, 73-78.
- Schroeder, A. N., & Best, T. M. (2015). Is self myofascial release an effective preexercise and recovery strategy? A literature review. *Current sports medicine reports*, 14(3), 200-208.
- Taşkuyu, E., & Özcan, S. (2025). Effect of different tennis court surfaces on the balance performance of tennis players. *Physical Education of Students*, 29(4), 277-283.
- Yamaguchi, T., & Ishii, K. (2005). Effects of static stretching for 30 seconds and dynamic stretching on leg extension power. *Journal of Strength and Conditioning Research*, 19(3), 677-683.
- Yamaguchi, T., Ishii, K., Yamanaka, M., & Yasuda, K. (2007). Acute effects of dynamic stretching exercise on power output during concentric dynamic constant external resistance leg extension. *Journal of Strength and Conditioning Research*, 21(4), 1238-1244.
-